

On the dielectric susceptibility calculation in the incommensurate phase of K_2SeO_4 .

T.A. Aslanyan*

Institute for Physical Research, Armenian National Academy of Sciences, Ashtarak-2, 378410 Armenia

It is shown that the thermodynamic potential of the domain-like incommensurate (IC) phase of the K_2SeO_4 crystal (viewed as a model for the IC-C transition) should be supplemented with a term, taking into account the local, Lorentz electric field. The latter qualitatively changes the result of calculation of the dielectric susceptibility for this IC structure by Nattermann and Trimper, J. Phys. C: Solid State Phys. **14**, 1603, (1981), and gives phase transition to the ferroelectric IC phase obtained by Aslanyan, Phys. Rev. B **70**, 024102, (2004).

PACS numbers: 64.70.Rh, 64.60.-i

In the present paper the behavior of the incommensurate (IC) phase of the K_2SeO_4 crystal near the lock-in transition is discussed. We remind that K_2SeO_4 undergoes on cooling two successive phase transitions: the IC transition at $T_i = 129K$ and the lock-in transition to the triple-period commensurate ferroelectric phase at $T_c = 93K$.^{1,2} Nattermann and Trimper³ calculated the dielectric susceptibility χ of the crystal in the IC phase, and showed that χ diverges on approaching the lock-in transition point (i.e., a loss of stability of the IC phase with respect to the commensurate phase takes place). The dielectric susceptibility for the same IC structure was calculated also by Aslanyan,⁴ and it was obtained that χ diverges not in the lock-in transition point, but at a higher temperature, giving rise to the phase transition from the IC to the ferroelectric IC structure. Spatial distributions of the IC modulation phase and polarization vector in the ferroelectric (or z -polarized) IC phase⁴ are depicted in Fig.1. Below we explain the origin of the difference between the results of the two calculations,^{3,4} and show that the thermodynamic potential used for calculations by Nattermann and Trimper³ should be supplemented with a term, taking into account the local (Lorentz) microscopic electric field, induced by the dipoles (the neighboring pairs of the domains in Fig.1 can be viewed as periodically arranged dipoles). In such a case, calculation of the dielectric susceptibility by Nattermann and Trimper³ should be revised, and it reduces to that carried out by Aslanyan.⁴

The starting point in both calculations^{3,4} is the thermodynamic potential of the following form:

$$\tilde{\Phi} = \int dx \left\{ \frac{D}{2} \eta_0^2 \left(\frac{\partial \varphi}{\partial x} \right)^2 + \frac{f}{2} \eta_0^6 \cos(6kx + 6\varphi) + \frac{r}{2} P \eta_0^3 \cos(3kx + 3\varphi) + \frac{\chi_0^{-1}}{2} P^2 \right\} \quad (1)$$

In this expression (and below as well) the notations of paper [4] are used, where η_0 and φ are, respectively, the amplitude and the phase of the IC modulation, k is the IC vector at the temperature of the IC transition (from the normal to the IC phase), P is the polarization vector in the z direction, and the IC modulation is given by $\eta_0 \cos(kx + \varphi)$.

However, potential (1) should be supplemented with a




FIG. 1: a) Coordinate x dependence of the z -polarized IC structure's modulation phase $u(x) = \varphi(x) + kx$. Each plateau corresponds to the domain with $+$ or $-$ polarization vector P , which is along the $\pm z$ direction (curve b). Since the sizes of the adjacent domains are different, the integral polarization of any two adjacent domains (and, consequently, of the crystal as a whole) is not zero. Polarization of the crystal appears due to the displacement of the odd domain walls with respect to the even walls, which makes nonequal the sizes of adjacent domains. The coordinate x is given in multiples of the lattice parameter. c) Periodic disposition of the dipoles in the $x - z$ plane. These dipoles correspond to the pairs of adjacent domains shown by the curve b). In the y and z directions the periods are equal to the crystal lattice parameters, while in the x direction the distance between dipoles equals to the width of the pair of domains.

term of the following origin. The domain-like IC structure, which develops from the sinusoidal one on cooling, can be polarized in the z direction via displacements of the odd domain walls with respect to the even walls in the x direction, as shown in Fig.1. The pairs of adjacent domains (which become non-equisize after such displacements) can be viewed as a periodically arranged dipoles as in Fig.1, c). However, any periodically (or chaotically, as in the case of a gas) distributed dipoles, with

nonzero integral polarization, induce a microscopic electric field E_{micr} (the so called local Lorentz field) biasing each dipole. For the case of a gas or a cubic crystal, this field is equal to $E_{micr} = 4\pi\bar{P}/3$, where $\bar{P}$ is the spatially averaged polarization. The effect of the microscopic electric field is implicitly taken into account for the susceptibility χ_0 of the high temperature phase. However, in the IC phase, when the crystal acquires a new sublattice in form of the polarizing pairs of domains, one should additionally take it explicitly into account. So, for the orthorhombic domain lattice of K_2SeO_4 the microscopic field is

$$E_{micr} = a\bar{P}, \text{ with } \bar{P} = \frac{1}{L} \int P(R)dx,$$

where a is a coefficient of the same order as $4\pi/3$ (in case of a cubic symmetry $a = 4\pi/3$), and L is the crystal's length in the x direction. Due to the appearance of the field, biasing each dipole, the thermodynamic potential (1) should be contributed by the additional term

$$-\bar{P}E_{micr} = -a\bar{P}^2 = -a \left(\int P(R)dx \right)^2 / L^2,$$

which should be added to the expression under integral (1). It should be emphasized a non-local character of this energy term (square of the integral), which significantly changes the expected results of calculations.

In what follows we show that the thermodynamic potential of the crystal with the domain-induced microscopic electric field can be reduced to the potential, analyzed in the paper [4], and the behavior of the system in such a case should be that described ibidem. Minimizing the potential over $P(R)$ and $\bar{P}$, under the condition

$$\int (P(r) - \bar{P})dx = 0,$$

which should be added to the potential equation with the Lagrange multiplier, one obtains

$$P(R) = -\frac{r\chi_0\eta_0^3}{2} \cos 3u - \frac{ar\chi_0^2\eta_0^3}{L(1-2\chi_0a)} \int \cos 3u dx$$

and

$$\bar{P} = -\frac{r\chi_0\eta_0^3}{2L(1-2a\chi_0)} \int \cos 3u dx,$$

where $u = kx + \varphi(x)$. One can substitute $P(R)$ and $\bar{P}$ back into the potential expression and obtain:

$$\begin{aligned} \tilde{\Phi} = & \int dx \left\{ \frac{D}{2} \eta_0^2 \left(\frac{\partial \varphi}{\partial x} \right)^2 + \frac{8f - r^2\chi_0}{16} \eta_0^6 \cos 6u - \right. \\ & \left. - \frac{r^2a\chi_0^2\eta_0^6}{4L^2(1-2a\chi_0)} \left(\int \cos 3u dx \right)^2 \right\} \end{aligned} \quad (2)$$

We once more note a non-local character of the coupling in this potential, which is given by the square of the integral in the equation. Such a coupling is induced by the

above-introduced microscopic electric field, and it disappears at $a = 0$. Eq.(2) allows one (via minimization over $\varphi(R)$) to derive the equation for the temperature evolving of the IC structure. For the case $a = 0$ the solution of the corresponding equation is well studied,⁵ and it gives IC domains of equal sizes, which are increasing (logarithmically diverging) on cooling towards the lock-in transition. The dielectric susceptibility for such a behavior was calculated by Nattermann and Trimper,³ who showed that it diverges at the lock-in transition point. However for the case $a \neq 0$, eq.(2) significantly differs from that for $a = 0$. It is easy to check that eq.(2) is equivalent to:

$$\begin{aligned} \tilde{\Phi} = & \int dx \left\{ \frac{D}{2} \eta_0^2 \left(\frac{\partial \varphi}{\partial x} \right)^2 + \frac{f_1}{2} \eta_0^6 \cos(6kx + 6\varphi) + \right. \\ & \left. \frac{r_1}{2} P_1 \eta_0^3 \cos(3kx + 3\varphi) + \frac{\chi_1^{-1}}{2} P_1^2 \right\}, \end{aligned} \quad (3)$$

where P_1 is a spatially constant vector and

$$\chi_1^{-1} = \chi_0^{-1} - 2a, \quad f_1 = f - r^2\chi_0/8, \quad r = r\sqrt{2a\chi_0}.$$

In order to check that at $a \neq 0$ eq.(3) should give the same solutions for the phase $\varphi(R)$ as eq.(2), it is sufficient to minimize eq.(3) over P_1 , and, as a result, return to eq.(2).

On the other hand, eq.(3) coincides by form with eq.(1), with P replaced by the spatially constant vector P_1 . Just the potential of eq.(3) with constant P_1 was used by Aslanyan⁴ (though the approach was not commented and sufficiently clarified there) for checking the stability of the IC phase, and it was shown that within the IC structure a phase transition should take place from the IC to the ferroelectric IC phase, prior to the lock-in transition on cooling. The spatial distribution of the phase $u = \varphi + kx$ and polarization $P(R)$ in this ferroelectric phase are those shown in Fig.1.⁴

In other words, for calculation of the dielectric susceptibility $\chi^{-1} = \partial^2 \tilde{\Phi} / \partial P^2$ at $P = 0$ one should minimize potential (1) with respect to φ , and find the solution of the equation:

$$D \frac{\partial^2 u}{\partial x^2} + 3f\eta_0^4 \sin 6u + \frac{3r}{2} P\eta_0 \sin 3u = 0.$$

In the paper ref.[4] the corresponding solution was found for the case $P = const$ in this equation (which corresponds to minimization of eq. (3)), and in ref.[3] that was obtained for a spatially dependent $P(R)$.

So, we showed that thermodynamic potential (1), used for calculation of the dielectric susceptibility, should be supplemented by the term, contributed from the domain-induced Lorentz local electric field. As a result, one can see that calculations of the dielectric susceptibility by Nattermann and Trimper³ give correct result only for a pair of the adjacent domains, which is separated from the crystal (i.e. for a single dipole). For a large number of pairs of domains, periodically disposed in space as in Fig.1 (in the real IC phase), the coupling between the

dipoles should be taken into account, which give phase transition to the ferroelectric IC phase with non-equisize IC domains. This coupling gives contribution of a negative sign to the potential, and makes ferroelectric IC phase more favorable compared to the equisize domain IC structure on approaching the lock-in transition.

As an experimental confirmation of the presented theory may be pointed out the second-harmonic generation in the IC phases of the K_2SeO_4 ⁶, $(\text{NH}_4)_2\text{BeF}_4$ ⁷ and, under some assumptions, of the quartz⁸ crystals, which is discussed in ref.[4], and the present paper should be

viewed as an additional foundation of its evidence. It is also worth to note that the developed theory allows one to explain such a surprising observations as a significant drop in the K_2SeO_4 elastic constant c_{55} (which corresponds to the transversal u_z sound propagating in the x direction) near the lock-in transition point⁹, and observation of a depolarized, overdamped Raman scattering in the $z(yz)x$ geometry¹⁰ in the IC phase. These observations unambiguously follow from the presented theory, and will be published elsewhere.

* Electronic address: aslanyan@freenet.am

¹ Y. Ishibashi, *Incommensurate Phases in Dielectrics*, edited by R. Blinc and A.P. Levanyuk, Modern Problems in Condensed Matter Sciences, Vol.**14.2**, North-Holland (1986).

² M. Iizumi, J.D. Axe, G. Shirane and K. Shimaoka, Phys. Rev. **B 15**, 4392 (1977).

³ T. Nattermann and S. Trimper, J. Phys. C: Solid State Phys., **14**, 1603, (1981).

⁴ T.A. Aslanyan, Phys. Rev. B **70**, 1, 024102 (2004)

⁵ I.E. Dzyaloshinski, Sov. Phys., JETP, **47**, 336, 992, (1964).

⁶ A.M. Arutyunyan, B. Brezina, S.Kh. Esayan, V.V. Le-

manov, Sov. Phys., Solid State, **24**, 814, (1982).

⁷ K.S. Alexandrov, A.N. Vtyurin, V.F. Shabanov, Sov. Phys., Pis'ma JETP, **28**, 153, (1978).

⁸ G. Dolino, J.P. Bachheimer, phys.stat.sol. (a), **41**, 673, (1977).

⁹ S.D. Prokhorova, G.A. Smolensky, I.G. Siny, E.G. Kuzimov, V.D. Mikvabia, H. Arndt, Ferroelectrics, **13**, 3823, (1980).

¹⁰ H.G. Unruh, W. Eller, G.Krit, phys.stat.sol. (a), **55**, 173, (1979).